
## Warm Up 4/9

Convert the following polar coordinates into rectangular coordinates.

1. 
$$(6, \frac{2\pi}{3}) = (6(0) \frac{2\pi}{3}, 6\sin \frac{2\pi}{3})$$
  
 $= (6(-1/2), 6(\frac{5\pi}{2}))$   
2.  $(0, 13\pi)$   
 $= (0(0)|3\pi, 0)\sin|3\pi$ 



### Objective

Students will...

Be able to convert rectangular equations into polar equations.

### Recap

So, for recap...

#### Relationship between Polar and Rectangular Coordinates

1. To change from polar to rectangular coordinates, use the formulas

$$x = r \cos \theta$$
 and  $y = r \sin \theta$ 

2. To change from rectangular to polar coordinates, use the formulas

$$r^2 = x^2 + y^2$$
 and  $\tan \theta = \frac{y}{x}$   $(x \neq 0)$ 

### **Polar Equations**

Polar Equations are equations solved for r, written in terms of  $\sin \theta$ ,  $\cos \theta$ , or a combination of both.

Since  $x = r \cos \theta$  and  $y = r \sin \theta$ , all rectangular equation (written in terms of x and y) can be written in polar equation form. We use the following steps.

Ex. Write the equation y = 2x - 9 in polar form.

Rewrite x as  $r\cos\theta$  and y as  $r\sin\theta$ :

implify and solve for r:

 $n\theta: \ rsin\theta = 2(r\cos\theta) - 9$   $rsin\theta = 2r\cos\theta - 9$   $2r\cos\theta - r\sin\theta = 9$   $r(2\cos\theta - \sin\theta) = 9$ 

### **Examples**

Express the equation 
$$x^2 = 4y$$
 in polar form.  $(V\cos\theta)^2 = 4(V\sin\theta)$ 

## Example

Convert the equation x = 1 to polar form.

#### Homework Problem

Convert the equation to polar form.

$$42. x^{2} + y^{2} = 9$$

$$(r \cos \theta)^{2} + (r \sin \theta)^{2} = 9$$

$$\Rightarrow r^{2}(\cos^{2}\theta + r^{2}\sin^{2}\theta) = 9$$

$$\Rightarrow r^{2}(\frac{\cos^{2}\theta + \sin^{2}\theta}{\cos^{2}\theta}) = 9$$

$$\Rightarrow r^{2}(\frac{\cos^{2}\theta + \sin^{2}\theta}{\cos^{2}\theta}) = 9$$

# Homework 4/9

TB pg. 587 #42-46