

### Objective

#### Students will...


- Be able to identify and solve for both vertical and horizontal asymptotes of rational functions.
- Be able to find the slant or oblique asymptote of a rational function, given that it exists.

# Asymptotes

One of the characteristics of rational function graphs is the presence of asymptotes. <u>Asymptotes</u> are lines that the graph of the function gets closer and closer to as it travels on. You may think of them as boundary lines that the graph continually approaches.

$$ex. f(x) = \frac{1}{x}$$

We can see that both x and the y-axis are asymptotes of this graph.



## **Vertical Asymptotes**

From the previous graph, we saw that there were two different types of asymptotes at play. There was a <u>vertical</u> asymptotes (the y-axis), as well as a <u>horizontal</u> asymptote (the x-axis). So for every rational function graph, we must consider both.

Recall that the vertical lines represent the horizontal or the x-coordinates. Thus, to find vertical asymptotes, we must consider the possible x-coordinates that would make the rational functions undefined, i.e. what x-value makes the denominator 0?

ex.  $f(x) = \frac{1}{x}$  For this function, it's obvious that the only place the function is undefined would be when x = 0, which is the y-axis. Therefore, it becomes the <u>vertical asymptotes</u>.

Find the vertical asymptotes of the following functions.

$$1. f(x) = \frac{x-6}{x+2}$$

$$2. g(x) = \frac{8}{2x - 9}$$

$$2x-9=0$$

$$2x = \frac{9}{9} = 4.5$$

$$3. h(x) = \underbrace{\frac{x-9}{5}}$$

VA: DNE

# **Horizontal Asymptotes**



Horizontal asymptotes are horizontal lines, which represents a certain y-value (y=...). The method for finding horizontal asymptotes is as follows:

Let n be the leading exponent of the numerator and m be the leading exponent of the denominator.

(a). If n < m, i.e. higher degree in the denominator, the horizontal asymptotes is y = 0.

(b). If n=m, then the horizontal asymptote is  $\frac{coefficient\ of\ leading\ term}{coefficient\ of\ leading\ term}$ 

(c). If n>m, i.e. higher degree in the numerator, then no horizontal asymptote exists

3×3-4

$$1. f(x) = \frac{x^2 - 6}{x^3 + 2}$$

Find the horizontal asymptotes of the following functions.


1. 
$$f(x) = \frac{x^2 - 6}{x^3 + 2}$$
2.  $g(x) = \frac{8k}{2k - 9}$ 
3.  $h(x) = \frac{9x^4}{5}$ 
4. A :  $y = \frac{9}{5} = \frac{4}{5}$ 
4. A :  $y = \frac{9}{5} = \frac{4}{5}$ 
4. S :  $y = \frac{9}{5} = \frac{4}{5}$ 

# Slant or Oblique Asymptotes

For situations where no horizontal asymptotes exist, i.e. higher degree in the numerator, there may still exist a <u>slant or oblique</u> asymptote. Finding such asymptote is a rather easy process, as it is simply done by dividing (long division is needed here).

Ex. 
$$f(x) = \frac{x^2-4x-5}{x^2-3} = > HA : DN &$$

Here we can easily see that no horizontal asymptote exists. We just need to divide to see if there is a slant or oblique asymptote.




Find the asymptotes of the following functions. If no horizontal asymptote exists, find slant or oblique asymptotes.

1. 
$$f(x) = \frac{5x+21}{x^2+10x+25}$$

VA:  $(x+5)(x+5) = 0$ 

VA:  $(x+5)(x+5) = 0$ 

VA:  $(x+5)(x+5) = 0$ 



$$3. f(x) = \frac{\int_{2x^2+4x}^{2-3x-4}}{2x^2+4x}$$

$$VA' \cdot Z \times {}^{2} + 4x = 0$$

$$2X(x+2) = 0$$

$$X = 0, -2$$

$$HA : \begin{cases} 4 - \frac{1}{2} \\ \frac{1}{2} \\ \frac{1}{2} \end{cases}$$

$$4. f(x) = \frac{2x^{2} + 7x - 4}{1)^{2} + x - 2}$$

$$(x + 1)(x - 1) = 0$$

$$(x + 1)(x - 1) = 0$$

$$(x - 2 - 1)$$

$$(x - 2 - 2)$$

$$(x - 2 - 2)$$

Homework 10/30

TB pg. 313 #11-23 (odd)